



## UWS Academic Portal

### **The toxicity of potentially toxic elements (Cu, Fe, Mn, Zn and Ni) to the cnidarian *Hydra attenuata* at environmentally relevant concentrations**

McKinley, Kevin; McLellan, Iain; Gagne, Francois; Quinn, Brian

*Published in:*  
Science of the Total Environment

*DOI:*  
[10.1016/j.scitotenv.2019.02.193](https://doi.org/10.1016/j.scitotenv.2019.02.193)

Published: 15/05/2019

*Document Version*  
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

#### *Citation for published version (APA):*

McKinley, K., McLellan, I., Gagne, F., & Quinn, B. (2019). The toxicity of potentially toxic elements (Cu, Fe, Mn, Zn and Ni) to the cnidarian *Hydra attenuata* at environmentally relevant concentrations. *Science of the Total Environment*, 665, 848-854. <https://doi.org/10.1016/j.scitotenv.2019.02.193>

#### **General rights**

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### **Take down policy**

If you believe that this document breaches copyright please contact [pure@uws.ac.uk](mailto:pure@uws.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# **The toxicity of potentially toxic elements (Cu, Fe, Mn, Zn and Ni) to the cnidarian *Hydra attenuata* at environmentally relevant concentrations**

Kevin McKinley <sup>a,\*</sup>, Iain McLellan <sup>a</sup>, Francois Gagné <sup>b</sup>, Brian Quinn <sup>a</sup>

<sup>a</sup> School of Health and Life Sciences, University of the West of Scotland, Paisley PA1 2BE, Scotland, United Kingdom

<sup>b</sup> Aquatic Contaminants Research Division, Environment and Climate Change Canada, 105 McGill street, Montréal, QC H2Y 2E7, Canada

## **1. Introduction**

Pollution of the aquatic environment is common near human activity and the presence of chromium (Cr), copper (Cu), manganese (Mn), nickel (Ni), lead (Pb), tin (Sn) and zinc (Zn) is associated with antifouling paint, industry, insecticides, fertilizer use, fuel consumption and waste water treatment works (Caccia *et al.*, 2003; Deheyn and Latz, 2006; Canning-Clode *et al.*, 2011; Xu *et al.*, 2014; Rodriguez-Iruretagoiena *et al.*, 2016). Areas with intense, localised activity (e.g. harbours and ports within estuaries) are known to exhibit greater anthropogenic influence (Birch *et al.*, 2015). An example of this is the Clyde Estuary, Scotland, which has since the Industrial Revolution received pollution from ship building, dye works and petroleum installations resulting it becoming the UK's most contaminated estuarine environment (Turner, 2000; Edgar *et al.*, 2003; Vane *et al.*, 2007; Vane *et al.*, 2011).

The Water Framework Directive (2000/60/EC) and Annex II of the Environmental Quality Standards Directive (2008/105/EC) include a number of PTEs (arsenic (As), cadmium (Cd), mercury (Hg), nickel (Ni)) which are classified as 'priority substances' and 'priority hazardous substances'. Requirements of the Water Framework Directive include a target of 'good' ecological status for waterbodies and

levels of pollutants within the Clyde have reduced over the last three decades and ecological recovery has been observed (Critchlow-Watton *et al.*, 2014).

Elements such as Cu, Fe, Zn and Mn, are essential in trace amounts to support and maintain functions in aquatic ecosystems (Tchounwou *et al.*, 2012), however Pb, Cu and Zn have previously found to be a 'triad' of metals associated with human influence, and at high concentrations are toxic, adversely impacting on human/animal health and the environment (ten Brink and Woudstra, 1991; McLellan *et al.*, 2013). To determine the effect of pollutants on aquatic organic substances, numerous studies exposing invertebrates to heavy metals have previously been undertaken (Lasier *et al.*, 2000; Borgmann *et al.*, 2005; Torres Guzmán *et al.*, 2010; García *et al.*, 2011; Liber *et al.*, 2011; Lopes *et al.*, 2014, Joško *et al.*, 2016). Karntanut and Pascoe (2002) exposed four different *Hydra* species (*vulgaris* (*Zurich*), *attenuata*, *oligactis* and *viridissima*) to varying concentrations of Cu, Cd and Zn. A variation in the lethal concentration (LC) between species was found with Cu being the most toxic of the three elements with an LC<sub>50</sub> ranging from 0.025 to 0.084 mg/l after 96h of exposure, followed by Cd (0.16 to 0.52 mg/l), then Zn (11 to 14 mg/l).

*H. attenuata* (also known as *Hydra vulgaris*) is a species of cnidarian that are ubiquitously found in freshwater ecosystems and are commonly used for toxicity testing. The health status and acute toxicity of *H. attenuata* is easily observed through a series of defined morphological changes following exposure to a toxin in a relatively simple bioassay (Wilby, 1988). Other chronic endpoints used to measure toxicity include asexual reproduction (budding), feeding behaviour and attachment to a substrate (Quinn *et al.*, 2012). This species has relatively unique regenerative properties, is easy to culture and maintain in a laboratory, has a high reproductive

rate and as a diploblastic organism, is sensitive to environmental pollutants and is therefore used as a bioindicator for the health of a freshwater aquatic ecosystem (Quinn *et al.*, 2012). *H. attenuata* have been widely used in cost effective bioassays to assess the toxicity of numerous contaminants including wastewater (Trottier *et al.*, 1997), industrial effluents (Blaise and Kusui, 1997), pharmaceuticals (Pascoe *et al.*, 2003; Quinn *et al.*, 2008a, 2008b, 2009) , PTEs (Holdway *et al.*, 2001; Karntanut and Pascoe, 2002; Quinn *et al.*, 2007) and more recently rare earth elements (Blaise *et al.*, 2018).

The aim of this study is to evaluate the toxicity of PTE's, both individually and as a mixture, found in the Clyde estuary in Scotland against the cnidarian *Hydra attenuata*. Water samples from various locations in the Clyde estuary (Scotland) were analysed for anthropogenic PTEs Cu, Fe, Mn, Ni and Zn. The toxicity of these metals individually and as a mixture at environmentally relevant and elevated concentrations was tested using the *H. attenuata* bioassay on the ecologically relevant endpoints of morphology, feeding, attachment and reproduction. To the best of our knowledge, this is the first toxicity study investigating the mortality of any *Hydra* species individually exposed to Fe, Mn or Ni.

## **2. Materials and Methods**

### **2.1. Test organism**

*Hydra* were maintained in glass bowls containing 0.5 L of *Hydra* media (147 mg/l CaCl<sub>2</sub>H<sub>2</sub>O, 110 mg/l TES [N-Tris(hydroxymethyl) methyl 1-2-aminoethanesulfonic acid], pH adjusted to 7 using 0.5 M NaOH), maintained at 18± 2°C with a 12 h light 12 h dark photoperiod. *Hydra* were fed 3 times per week with newly hatched *Artemia*

*salina* nauplii and were fasted 48 h prior to exposure. To avoid algal contamination, *Hydra* media was regularly changed after each feeding.

## **2.2. PTE determination**

Water samples were collected from two estuarine (Kelburn Park and Erskine Harbour) and one freshwater (Gourock Burn) sites along the Clyde estuary (Fig. 1). Samples were collected in polypropylene sample bottles with pH and temperature recorded immediately (Mettler Toledo). Samples were then acidified with conc. HNO<sub>3</sub> (Fisher Trace Grade, UK) on site for preservation, transported to the laboratory and refrigerated at 4°C until analysis. Prior to analysis samples were filtered to <45µm (Filtermate, Environmental Express, USA).

Potentially toxic elements were determined by ICP-OES (Thermo Fisher, iCAP); a calibration series (0 mg/l, 2 mg/l and 10 mg/l of multi-element standard, ME/1001/05; Fisher Scientific, UK) was determined. Samples were analysed in triplicate. ICP-OES conditions were as follows: rf generator: 1.15 kW; Plasma: 1.4 l/min; Auxillary: 0.5 l/min; Nebuliser: 0.8 l/min; sample flow rate 1.5 ml/min.

Averages of the sample concentrations were calculated; Limits of Detection were calculated using standard practice (e.g. (McLellan *et al.*, 2013)) (Table 1). It can be seen that levels of Fe within the Gourock Burn were very high therefore it was decided not to put this forward at the reported concentrations. Ni was taken at 0.5 mg/l to reflect potential toxicity levels within the selected biota. These are the 'environmentally relevant' concentrations.

### 2.3. Test solutions

Environmental relevant PTE solutions (1x) of metals in hydra media (HM) were prepared for the individual elements and for the mix solution. These stock solutions were then diluted with HM to give concentrations: 0.0001x, 0.001x, 0.01x, 0.1x, 10x, 100x and 1000x concentrations were made from a 1000x stock solution (Table 2).

### 2.4. *Hydra* toxicity tests

All PTE exposures to *Hydra attenuata* were undertaken in quadruplicate (4 repetitions of each concentration) and the whole experiment was undertaken in triplicate for the PTE mixture and duplicate for each individual metal exposure. A 4 ml sample of the relevant solution was added to 4 wells in a 24 multiwell plate, containing a single *Hydra*, and the wells wrapped in parafilm to prevent evaporation and kept at  $18 \pm 2^\circ\text{C}$  for 96 h. Healthy *Hydra* with a morphology score of 10 on the Wilby table (Table 3) and having one bud (2 hydranths) were used in each exposure. Selection of healthy *Hydra* was undertaken using a binocular microscope. Morphology, hydranth number and attachment were observed at 24, 48, 72 and 96 h. The *Hydra*'s ability to ingest prey (feeding endpoint) was tested on all *Hydra* that scored  $> 5$  on the Wilby score table after 96 h as per Quinn et al., (2007). These *Hydra* were placed individually into a well of a clean 24 well multi-well plate containing 4 ml of *Hydra* media. Freshly hatched *Artemia* were rinsed three times with HM with 5 individuals added to each well at time 0, taking care not to add them directly to the tentacles of the *Hydra*. The number of ingested prey were observed every 20 min for 120 min.

## 2.5. Statistical Data Analysis

The 96 h LC<sub>50</sub> values for the mortality exposure were calculated using the Probit analysis program. The mortality exposure the sub-lethal LOEC (Lowest Observable Effect Concentration) was reported for  $\geq 2$  *Hydra* with score 8 or below and the NOEC (No Observable Effect Concentration) was based on *Hydra* with a score  $>8$  (Quinn et al., 2009). A toxicity threshold (TT) was determined from the LOEC and NOEC using the following equation:  $TT = (NOEC \times LOEC) / 2$  (US EPA, 1989). Variability in all endpoints (morphology, attachment, hydranth number, feeding behaviour) between the exposed and control *Hydra* were tested by one-way analysis of variance (ANOVA). Significance was set at  $p \leq 0.05$ . The Pearson correlation coefficient was used to measure the strength of the association between the concentration of the pollutant and the endpoints.

## 3 Results

### 3.1. Toxicity of individual metals to *Hydra attenuata*

Complete (100%) population mortality (indicated by a score  $\leq 5$  on the Wilby scale) was found at 0.1x (0.05 mg/l) for Cu (Fig. 2 A), 0.1x (0.3 mg/l) for Fe (Fig. 2 C), 10x (5 mg/l) for Ni (Fig. 2 D). For Mn, 100% mortality of all *Hydra* exposed was found at 100x (200 mg/l). Although some mortality was observed at 1x, mortality numbers were low (Fig. 2 B). Highly significant ( $p = < 0.005$ ) and negative correlations were found with hydranth number and feeding behaviour (Table 5). For Zn, 100% mortality was found at 1000x (100 mg/l). Although mortality was detected at 100x (10 mg/l), mortality numbers were low (Fig. 2 E). An extremely significant ( $p = < 0.001$ ) and

negative correlation was found with hydranth number, and a very significant ( $p = < 0.005$ ) negative correlation was found for feeding behaviour (Table 5). The 96 h  $LC_{50}$  values were determined as follows: Cu 0.0225 mg/l, Mn 20 mg/l, Fe 0.135 mg/l, Ni 2.25 mg/l, and Zn 31.622 mg/l. The Toxicity thresholds were calculated at: Cu 0.000125 mg/l, Mn 0.2 mg/l, Fe 0.000045 mg/l, Ni 0.0125 mg/l, Zn 5 mg/l (Table 4).

### 3.2 Toxicity of PTE mixture

For the PTE mixture, 100% mortality was found at 0.1x (Fig. 3). The 96h  $LC_{50}$  value was calculated as 0.045x. The LOEC was 0.01x and NOEC was 0.001x. The toxicity threshold was calculated at 0.000005x (Table 4). The high toxicity of Cu was not entirely responsible for the very high toxicity of the mixture. The toxicity threshold for the mixture (0.000005x) showed that the mixture was more toxic than Cu individually, which had a toxicity threshold of 0.000125 mg/l (0.00025x).

### 3.3 Toxicity of heavy metals at environmental concentration

Both Cu and Fe when exposed individually to the concentration of their respective metals found in the environment resulted in 100% mortality of all *Hydra* exposed (Fig. 2 A & C). A significant ( $p = < 0.001$ ) toxic effect occurred when *Hydra* were exposed to Mn and Ni at the environmentally relevant concentration. Zn remained at a perfect morphology score of 10 when exposed to the Zn concentration found in the environment (Fig. 2 E). The concentration of Zn found in the environment also had no significant toxic effect on hydranth number, feeding behaviour or attachment of *Hydra* to a substrate. When exposed to the concentration of Mn found in the environment, a significant ( $p = < 0.01$ ) toxic effect occurred in the attachment of *Hydra* to a substrate (Fig. 2 B). The concentration of Mn found in the environment had no significant toxic



effect on hydranth number or feeding behaviour. The concentration of Ni found in the environment resulted in a significant ( $p = < 0.05$ ) toxic effect on the feeding behaviour of *Hydra* (Fig. 2 D). The concentration of Ni found in the environment had no significant toxic effect on hydranth number or attachment of *Hydra* to a substrate. *Hydra* morphology was monitored at 24 h, 48 h, 72 h and 96 h of exposure to the concentration found in the environment with any toxic effect occurring within the first 24 h of exposure (Fig. 4).

#### 4 Discussion

Since the 18<sup>th</sup> Century and the beginning of the Industrial Revolution, the Clyde has had a diverse heritage and there is a well-documented legacy of pollutants e.g. (Hursthouse *et al.*, 1994; Edgar *et al.*, 2003; Vane *et al.*, 2007; Vane *et al.*, 2011). The sample locations chosen for this site are near former landfill sites (Gourock Burn and Kelburn Park) or wastewater treatment works (Erskine Harbour) and there is potential for continued contamination from these sources. This is in addition to former industrial activity e.g. metal plating near Kelburn Park (Miller, 1986). Despite the improving physical and ecological status of the outer Clyde estuary (Critchlow-Watton *et al.*, 2014), it is concerning that this study has found that PTE levels are above legislative requirements (Table 6) which may be caused by the proximity of potential point sources of pollutants. In that light, the Clyde is similar to other estuaries where point sources can be attributed to elevated PTE levels (Larrose *et al.*, 2010; Birch *et al.*, 2015; Petit Jérôme *et al.*, 2015; Rodriguez-Iruretagoiena *et al.*, 2016). Levels of all heavy metals tested were higher than levels in the Thames river in London, Canada (Environment and Engineering Services, 2018) and the Ganga river in India (Central Water Commission, 2018) (Table 6). The maximum acceptable

limits for copper (0.00376 mg/l) and iron (1 mg/l) based on EU / UK legislative requirements are higher than the *H. attenuata* LC<sub>50</sub>'s for copper (0.0225 mg/l) and iron (0.135 mg/l).

To the best of our knowledge, this is the first toxicity study investigating the toxicity of any *Hydra* species exposed to Fe, Mn or Ni. This study calculated the LC<sub>50</sub> values, LOEC, NOEC and Toxicity Thresholds for Cu, Fe, Mn, Zn and Ni (Table 4). The 96 h LC<sub>50</sub> results for Cu (0.0225mg/l) are similar to those reported by Karntanut and Pascoe (2000) (0.032 mg/l) for *H. vulgaris* (also known as *H. attenuata*) and for 4 different species of *Hydra*; *H. vulgaris* Zurich (0.042 mg/l), *H. vulgaris* (0.056 mg/l), *H. oligactis* (0.084 mg/l), *H. viridissima* (0.025 mg/l) (Karntanut and Pascoe, 2002). The Cu LC<sub>50</sub> value in the current study were higher than the LOEC value which is unusual but is due to the dilution range used for the serial dilution.

The 96 h LC<sub>50</sub> value calculated for Zn in the present study (31.6 mg/l) is higher than those reported for *H. vulgaris* (7.4 mg/l) (Karntanut & Pascoe, 2000) *H. vulgaris* Zurich (14 mg/l), *H. vulgaris* (13 mg/l), *H. oligactis* (14 mg/l), *H. viridissima* (11 mg/l) (Karntanut & Pascoe, (2002). In the current study Zn was tested at a concentration of 10 mg/l and a mortality percentage of 12.5% was found. The large divisions used in the serial dilutions resulted the high LC<sub>50</sub> value of 30 mg/l that was calculated, as the next concentration tested after 10 mg/l was 100 mg/l.

The same could be true of Mn (with an LC<sub>50</sub> value of 20 mg/l) but as this is the first time this metal has been used in a toxicity test to study mortality of *Hydra*, there is no literature for comparison. Harford *et al.*, (2015) however, exposed *Hydra viridissima* to varying levels of Mn to test population growth. The highest concentration tested by Harford *et al.*, (2015) was 10 mg/l at which the population of

*H. viridissima* was still growing but had dropped to 10% growth compared to the control. A very significant ( $p = <0.005$ ) negative correlation was found for hydranth number and feeding behavior when exposed to Mn.

An extremely significant ( $p = <0.001$ ) negative correlation for hydranth number and a very significant ( $p = <0.005$ ) negative correlation was found when exposed to Zn. However, there was no significant correlation for attachment when exposed to any of the tested metals and no significant correlation for attachment, hydranth number or feeding behavior when exposed to Cu, Fe or Ni.

In this study, a significant toxic effect occurred when *Hydra* were exposed to the Cu, Fe, Mn and Ni at concentrations found in the Clyde estuary (Fig. 2A-D). *Hydra* morphology was unaffected and remained at a score of 10 when exposed to the concentration of Zn found in the environment (Fig. 2E). Mortality levels of 100% were measured when *Hydra* were exposed to the heavy metal mixture (Fig. 3) and to Cu and Fe (individually) (Fig. 2A, C) at concentrations found in the Clyde. These results indicate that *Hydra attenuata* are unable to survive in aquatic environments with the metal concentrations found in the Clyde estuary, which may also have an impact on *Hydra* predators and prey.

The results also indicate that the PTE mixture (including the individual concentrations of Cu, Fe, Mn and Ni) could potentially prove significantly toxic to other invertebrates. The concentration of Cu found in the Clyde estuary was measured at 0.5 mg/l, this was 22 times higher than the  $LC_{50}$  found for *Hydra attenuata*. When compared with other studies (Table 7), the levels of Cu found in the Clyde would also be toxic to aquatic vertebrates such as *Rasbora sumatrana*, the guppy (*Poecilia reticulata*) and the zebrafish (*Danio rerio*). The concentration of Fe

found in the Clyde estuary was measured at 3 mg/l, this was also 22 times higher than the LC<sub>50</sub> found for *Hydra attenuata* and would be toxic to other aquatic invertebrates such as *Daphnia magna*, and aquatic vertebrates, such as the brown trout (*Salmo trutta*) (Table 7).

For the PTE mixture, a significant ( $p \leq 0.05$ ) toxic effect was seen at the lowest concentration studied (0.0001x) (Fig. 3). The LC<sub>50</sub> was calculated as 0.045x for the mixture and the toxicity threshold was calculated as 0.000005x. The toxicity threshold (TT) was lower than any of the corresponding values of the individual metals contained within the mixture (Table 4). This result indicates that the metals have a cumulative effect, with each metal behaving cumulatively, contributing to the total effect of the mixture and further increasing the toxicity.

Morphology was found to be the most significant endpoint in studying the toxic effects of metals. Using the additional endpoints of hydranth number, attachment and feeding behavior, Quinn *et al.*, (2007) found a significant decrease in hydranth number, attachment and feeding behavior as the concentration of the toxin increased. In the present study, a similar significant negative correlation was observed for hydranth number and feeding behavior following exposure to Mn and Zn. There was no significant correlation found with attachment in any of the exposures undertaken.

Most toxicity tests involving *Hydra* spp expose the organism to a toxin for 96 h. In this study, it was observed that any significant toxic effect of a pollutant occurred within the first 24 h of exposure. A review of other toxicity studies using *Hydra* as a test organism shows that the toxic effect of a contaminant occurs within the first 24 h of exposure (Blaise and Kusui, 1997; Karntanut and Pascoe, 2000, 2002). It may

therefore be necessary to only expose *Hydra* to a toxin for 24 h to test a compounds toxicity. However, more research is needed to confirm this. The potential replacement of a 96 h exposure with a 24 h one would greatly reduce the time needed for toxicity testing, helping to reduce the cost and potentially increasing the number of toxins that can be tested within a given time period.

## 5. Conclusion

This paper shows that a significant toxic effect was observed on *Hydra* exposed to the PTE mixture at the concentration found in the environment (1x) after a short-term exposure period (24 h). The high toxicity of Cu was not entirely responsible for the very high toxicity of the mixture. The toxicity threshold for the mixture (0.000005x) showed that the mixture was more toxic than Cu individually, which had a toxicity threshold of 0.000125 mg/l (0.00025x). The toxicity threshold (TT) for the PTE mixture was lower than that for the same metals when exposed individually to *Hydra*, indicating that metals may act cumulatively in a mixture. However, a significant toxic effect occurred when *Hydra* were exposed individually to Cu, Fe, Mn and Ni at concentrations found in the environment, with 100% mortality when exposed individually to the environmental concentrations of Cu and Fe. These high environmental concentrations of PTE would impact, not only on the predator and prey interactions within the *Hydra* community but also could potentially prove significantly toxic to other aquatic organisms.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## References

Alsop, D. and Wood, C. (2011) "Metal uptake and acute toxicity in zebrafish: common mechanisms across multiple metals." *Aquatic Toxicology*, 105 (3-4), pp. 385-393

Birch, G. F., Gunns, T. J. and Olmos, M. (2015) "Sediment-bound metals as indicators of anthropogenic change in estuarine environments." *Marine Pollution Bulletin*, 101(1), 243-257.

Blaise, C. and Kusui, T. (1997) 'Acute toxicity assessment of industrial effluents with a microplate-based *Hydra attenuata* assay', *Environmental Toxicology and Water Quality*, 12(1), pp. 53–60.

Blaise, C. Gagné, F. Harwood, M. Quinn, B. Hanana, H. (2018) 'Ecotoxicity responses of the freshwater cnidarian *Hydra attenuata* to 11 rare earth elements', *Ecotoxicology and Environmental Safety*, 163, pp. 486–491.

Borgmann, U. Couillard, Y. Doyle, P. Dixon, D. (2005) 'Toxicity of sixty-three metals and metalloids to *Hyalella azteca* at two levels of water hardness', *Environmental Toxicology and Chemistry*, 24(3), p. 641.

Caccia, V. G., Millero, F. J. and Palanques, A. (2003) "The distribution of trace metals in Florida Bay sediments." *Marine Pollution Bulletin* 46(11): 1420-1433.

Canning-Clode, J. Fofonoff, P. Riedel, G. F. Torchin, M. Gregory M. (2011) 'The Effects of Copper Pollution on Fouling Assemblage Diversity: A Tropical-Temperate Comparison', *PLoS ONE*, 6(3), p. e18026.

321 Central Water Commission (2018). *Status of Trace and Toxic Metals in Indian Rivers*.  
 322 Government of India, pp.1-251.

323 Critchlow-Watton, N., Dobbie, K., Bell, R., Campbell, S. G., Hinze, D., Motion, A.,  
 324 Robertson, K., Russell, M., Simpson, J., Thomson, D. and Towers, W. (2014)  
 325 "Scotland's State of the Environment Report." Scotland's Environment Web.

326 Dalzell, D. J. B. and MacFarlane N. A. A. (1999) "The toxicity of iron to brown trout and  
 327 effects on the gills: a comparison of two grades of iron sulphate" *Journal of Fish*  
 328 *Biology*, 55 (2), pp. 301-315

329 Deheyn, D. D. and Latz, M. I. (2006) 'Bioavailability of metals along a contamination  
 330 gradient in San Diego Bay (California, USA)', *Chemosphere*, 63(5), pp. 818–834.

331 Ebrahimpour, M. Alipour, H. Rakhshah, S. (2010) "Influence of water hardness on  
 332 acute toxicity of copper and zinc on fish." *Toxicology and Industrial Health*, 26 (6), pp.  
 333 361-365

334 Edgar, P. J., Hurthouse, A. S., Matthews, J. E. and Davies, I. M. (2003) "An  
 335 investigation of geochemical factors controlling the distribution of PCBs in intertidal  
 336 sediments at a contamination hot spot, the Clyde Estuary, UK." *Applied Geochemistry*  
 337 18(2): 327-338.

338 Environment and Engineering Services (2018). *Thames River Water Quality 2017*.  
 339 City of London, pp.1-10.

340 García, A. Espinosa, R. Delgado, L. Casals, E. González, E. Puntos, V. Barata, C.  
 341 Font, X. Sánchez, A. (2011) 'Acute toxicity of cerium oxide, titanium oxide and iron  
 342 oxide nanoparticles using standardized tests', *Desalination*, 269(1–3), pp. 136–141.

343 Harford, A. J. Mooney, T. J. Trenfield, M. A. van Dam, R. A. (2015) 'Manganese  
 344 toxicity to tropical freshwater species in low hardness water', *Environmental*  
 345 *Toxicology and Chemistry*, 34(12), pp. 2856–2863.

346 Hoseini, S. M. Rajabiesterabadi, H. Kordrostami, S. (2014) "Chronic exposure of  
 347 *Rutilus rutilus caspicus* fingerlings to ambient copper: Effects on food intake, growth  
 348 performance, biochemistry and stress resistance." *Toxicology and Industrial Health*,  
 349 32 (2), pp. 375-383

350 Holdway, D. A., Lok, K. and Semaan, M. (2001) 'The acute and chronic toxicity of  
 351 cadmium and zinc to two hydra species.', *Environmental toxicology*, 16(6), pp. 557-  
 352 65.

353 Hurthouse, A. S., Adamczyk, M., Adamczyk, L., Smith, F. J. and Iqbal, P. (1994)  
 354 "Inorganic and organic contaminants in intertidal sediments of the Clyde: preliminary  
 355 observations of historic trends." *Marine Pollution Bulletin* 28(12): 765-767.

356 Joško, I., Oleszczuk, P., Skwarek, E. (2016) 'The bioavailability and toxicity of ZnO and  
 357 Ni nanoparticles and their bulk counterparts in different sediments' *Journal of Soils and*  
 358 *Sediments*, 16(6), pp. 1798-1808

359 Karntanut, W. and Pascoe, D. (2000) 'A comparison of methods for measuring acute  
 360 toxicity to *Hydra vulgaris*', *Chemosphere*, 41(10), pp. 1543–1548.

361 Karntanut, W. and Pascoe, D. (2002) 'The toxicity of copper, cadmium and zinc to  
 362 four different *Hydra* (Cnidaria: Hydrozoa)', *Chemosphere*, 47, pp. 1059–1064.

363 Larrose, A., Coynel, A., Schäfer, J., Blanc., Massé, L. and Maneux, E. (2010)  
 364 "Assessing the current state of the Gironde Estuary by mapping priority contaminant  
 365 distribution and risk potential in surface sediment." *Applied Geochemistry* 25(12):  
 366 1912-1923.



367 Lasier, P. J., Winger, P. V. and Bogenrieder, K. J. (2000) 'Toxicity of Manganese to  
 368 *Ceriodaphnia dubia* and *Hyalella azteca*', *Archives of Environmental Contamination*  
 369 *and Toxicology*, 38(3), pp. 298–304.

370 Liber, K., Doig, L. E. and White-Sobey, S. L. (2011) 'Toxicity of uranium,  
 371 molybdenum, nickel, and arsenic to *Hyalella azteca* and *Chironomus dilutus* in water-  
 372 only and spiked-sediment toxicity tests', *Ecotoxicology and Environmental Safety*,  
 373 74(5), pp. 1171–1179.

374 Lopes, S. Ribeiro, F. Wojnarowicz, J. Łojkowski, W. Jurkschat, K. Crossley, A.  
 375 Soares, A. M. V. M. Loureiro, S. (2014) 'Zinc oxide nanoparticles toxicity to *Daphnia*  
 376 *magna*: size-dependent effects and dissolution', *Environmental Toxicology and*  
 377 *Chemistry*, 33(1), pp. 190–198.

378 McLellan, I., Hurthouse, A., Varela, A. and Pereira, C. S. (2013) "Geochemical  
 379 approach to assessing human impacts in Cork Oak forest soils of the MED region."  
 380 *Journal of Geochemical Exploration* 132: 34-40.

381 Miller, B. S. (1986) 'Trace metals in the common mussel *Mytilus edulis* (L.) in the  
 382 Clyde Estuary', *Proceedings of the Royal Society of Edinburgh*, 90, pp. 377–391.

383 Ololade, I. A. and Oginni, O. (2010) "Toxic Stress and Hematological Effects of  
 384 Nickel on African Catfish, *Clarias gariepinus*, Fingerlings." *Journal of Environmental*  
 385 *Chemistry and Ecotoxicology*, 2, pp. 14-19

386 Pascoe, D., Karntanut, W. and Müller, C. T. (2003) 'Do pharmaceuticals affect  
 387 freshwater invertebrates? A study with the cnidarian *Hydra vulgaris*', *Chemosphere*,  
 388 51(6), pp. 521–528.

389 Petit Jérôme, C. J., Schäfer, J., Coynel, A., Blanc, G., Chiffolleau, J. F., Auger, D.,  
 390 Bossy, C., Derrienic, H., Mikolaczyk, M., Dutruch, L. and Mattielli, N. (2015) "The  
 391 estuarine geochemical reactivity of Zn isotopes and its relevance for the biomonitoring  
 392 of anthropogenic Zn and Cd contaminations from metallurgical activities: Example of  
 393 the Gironde fluvial-estuarine system, France." *Geochimica et Cosmochimica Acta* 170:  
 394 108-125.

395 Quinn, B., Gagné, F. and Blaise, C. (2007) 'Validation of a multi-well plate toxicity  
 396 test to assess feeding behaviour of the cnidarian, *Hydra attenuata*', *Fresenius*  
 397 *Environmental Bulletin*, 16(9 A), pp. 1100–1107.

398 Quinn, B., Gagné, F. and Blaise, C. (2008a) 'An investigation into the acute and  
 399 chronic toxicity of eleven pharmaceuticals (and their solvents) found in wastewater  
 400 effluent on the cnidarian, *Hydra attenuata*', *Science of the Total Environment*, 389(2–  
 401 3), pp. 306–314.

402 Quinn, B., Gagné, F. and Blaise, C. (2008b) 'The effects of pharmaceuticals on the  
 403 regeneration of the cnidarian, *Hydra attenuata*', *Science of the Total Environment*,  
 404 402(1), pp. 62–69.

405 Quinn, B., Gagné, F. and Blaise, C. (2009) 'Evaluation of the acute, chronic and  
 406 teratogenic effects of a mixture of eleven pharmaceuticals on the cnidarian, *Hydra*  
 407 *attenuata*', *Science of the Total Environment*, 407(3), pp. 1072–1079.

408 Quinn, B., Gagné, F. and Blaise, C. (2012) '*Hydra*, a model system for environmental  
 409 studies', *The International Journal of Developmental Biology*, 56(6-7–8), pp. 613–  
 410 625.

411 Rodriguez-Iruretagoiena, A., elejoste, N., Gredilla, A., Fdez-Ortiz de Vallejuelo, S.,  
 412 Arana, G., Madariaga, J.M. and de Diego, A. (2016) "Occurrence and geographical  
 413 distribution of metals and metalloids in sediments of the Nerbioi-Ibaizabal estuary  
 414 (Bilbao, Basque Country)." *Marine Chemistry* 185: 82-90.

415 Scottish Environment Protection Agency (2018) "Supporting Guidance (WAT-SG-53):  
 416 Environmental Quality Standards for Discharges to Surface Waters"

417 Sharma, J. Langer, S. (2014) "Effect of Manganese on haematological parameters of  
 418 fish, *Garra gotyla gotyla*" *Journal of Entomology and Zoology Studies*, 2 (3), pp. 77-  
 419 81

420 Shuhaimi-Othman, M. Nadzifah, Y. and Ahmad, A.K. (2010) "Toxicity of copper and  
 421 cadmium to freshwater fishes" *World Academy of Science, Engineering and*  
 422 *Technology*, 65, pp. 869-871

423 ten Brink, B. J. E. and Woudstra, J. H. (1991). "Towards an effective and rational  
 424 water management: the aquatic outlook project-integrating water management,  
 425 monitoring and research." *European Water Pollution Control*, 1, pp. 20-27

426 Tchounwou, P. B. Yedjou, C. G. Patlolla, A. K. Sutton, D. J. (2012) 'Heavy metal  
 427 toxicity and the environment.', *EXS. NIH Public Access*, 101, pp. 133–64.

428 Torres Guzmán, F., González, F. J. A. and Rico Martínez, R. (2010) 'Implementing  
 429 *Lecane quadridentata* acute toxicity tests to assess the toxic effects of selected  
 430 metals (Al, Fe and Zn)', *Ecotoxicology and Environmental Safety*, 73(3), pp. 287–  
 431 295.

432 Trottier, S. Blaise, C. Kusui, T. Johnson, E. M. (1997) 'Acute toxicity assessment of  
 433 aqueous samples using a microplate-based *Hydra attenuata* assay', *Environmental*  
 434 *Toxicology and Water Quality*, 12(3), pp. 265–271.

435 Turner, A. (2000) 'Trace Metal Contamination in Sediments from U.K. Estuaries: An  
 436 Empirical Evaluation of the Role of Hydrous Iron and Manganese Oxides', *Estuarine,*  
 437 *Coastal and Shelf Science*, 50(3), pp. 355–371.

438 Vane, C., Harrison, I. and Kim, A. W. (2007) "Assessment of polyaromatic  
 439 hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) in surface sediments of  
 440 the Inner Clyde Estuary, UK." *Marine Pollution Bulletin* 54(8): 1301-1306.

441 Vane, C., Chenery, S. R., Harrison, I., Kim, A. W., Moss-Hayes, V. and Jones, D. G.  
 442 (2011) "Chemical signatures of the Anthropocene in the Clyde estuary, UK: sediment-  
 443 hosted Pb, <sup>207/206</sup>Pb, total petroleum hydrocarbon, polyaromatic hydrocarbon and  
 444 polychlorinated biphenyl pollution records." *Philosophical transactions of the Royal*  
 445 *Society A* 369: 1085-1111.

446 Wilby, O. K. (1988). The *Hydra* regeneration assay. *Proceedings of Workshop*  
 447 *organised by Association Francaise de Teratologie*: 108-124.

448 Xu, X. Zhao, Y. Zhao, X. Wang, Y. Deng, W. (2014) 'Sources of heavy metal  
 449 pollution in agricultural soils of a rapidly industrializing area in the Yangtze Delta of  
 450 China', *Ecotoxicology and Environmental Safety*, 108, pp. 161–167.


451

452

453

454

455



456

457 Fig. 1: Overview of heavy metal sampling locations along the Clyde estuary, Scotland. \* indicates  
458 sample location.

459

460





Fig. 2: – Lethal and sub-lethal effects of individual metals at varying concentrations on *Hydra* morphology, hydranth number, attachment and feeding after 96hr exposure. Points at Morphology and feeding represent the mean score ( $n=8$ ) $\pm$ standard error. Points at attachment and hydranth number represent sum( $n=8$ ). Significance for morphology at \*= $p\leq 0.05$ ; \*\*= $p\leq 0.01$ ; \*\*\*= $p\leq 0.001$ . Significance for hydranth number at  $\blacksquare$ = $p\leq 0.05$ ;  $\blacksquare\blacksquare$ = $p\leq 0.01$ ;  $\blacksquare\blacksquare\blacksquare$ = $p\leq 0.001$ . Significance for attachment at  $\blacktriangle$ = $p\leq 0.05$ ;  $\blacktriangle\blacktriangle$ = $p\leq 0.01$ ;  $\blacktriangle\blacktriangle\blacktriangle$ = $p\leq 0.001$ . Significance for feeding at  $\blacktriangledown$ = $p\leq 0.05$ ;  $\blacktriangledown\blacktriangledown$ = $p\leq 0.01$ ;  $\blacktriangledown\blacktriangledown\blacktriangledown$ = $p\leq 0.001$ . Note: Error bars do not show at points where results had no variability. (Should be printed in colour)



474

475 Fig. 3 – Lethal and sub-lethal effects of several concentrations (0.0001x–1x) of the heavy metal  
 476 mixture found in the environment on *Hydra* morphology, hydranth number, attachment and feeding  
 477 after 96hr exposure. Points at Morphology and feeding represent the mean score (n=12)  $\pm$ standard  
 478 error. Points at attachment and hydranth number represent sum (n=12). Significance for morphology  
 479 at \*= $p \leq 0.05$ ; \*\*= $p \leq 0.01$ ; \*\*\*= $p \leq 0.001$ . Significance for hydranth number at  $\blacksquare$ = $p \leq 0.05$ ;  $\blacksquare$ = $p \leq 0.01$ ;  
 480  $\blacksquare$ = $p \leq 0.001$ . Significance for attachment at  $\blacktriangle$ = $p \leq 0.05$ ;  $\blacktriangle$ = $p \leq 0.01$ ;  $\blacktriangle$ = $p \leq 0.001$ . Significance for  
 481 feeding at  $\blacktriangledown$ = $p \leq 0.05$ ;  $\blacktriangledown$ = $p \leq 0.01$ ;  $\blacktriangledown$ = $p \leq 0.001$ .

482

483

484




Fig. 4 – Effect of heavy metal concentrations found in the environment on *Hydra* morphology at 24, 48, 72 and 96 h exposure. Mean scores are represented the individual metals (n=8) and metal mixture (n=12)±standard error. Significance for morphology at \*=p≤0.05; \*\*=p≤0.01; \*\*\*=p≤0.001. Significance for Cu at ■=p≤0.05; ■■=p≤0.01; ■■■=p≤0.001. Significance for Fe at ▲=p≤0.05; ▲▲=p≤0.01; ▲▲▲=p≤0.001. Significance for Mn at ▼=p≤0.05; ▼▼=p≤0.01; ▼▼▼=p≤0.001. Significance for Ni at ▲=p≤0.05; ▲▲=p≤0.01; ▲▲▲=p≤0.001.

(Should be printed in colour)

Table 1: Estuarine and freshwater concentrations of heavy metals found in the environment.

| Element   | Gourock Burn (mg/l) | Kelburn Park (mg/l) | Erskine Harbour (mg/l) | Average (mg/l) |
|-----------|---------------------|---------------------|------------------------|----------------|
| <b>Cu</b> | <LOD                | 0.98                | 0.67                   | 0.82           |
| <b>Fe</b> | 33.78               | <LOD                | 9.87                   | 21.82          |
| <b>Mn</b> | 1.72                | <LOD                | <LOD                   | 1.72           |
| <b>Ni</b> | <LOD                | 1.74                | 1.42                   | 1.58           |
| <b>Zn</b> | <LOD                | 0.20                | 0.23                   | 0.21           |

Table 2: The concentrations of PTEs used in the exposure tests (mg/l). Based on the concentrations found in the environment.

| Metal            | 0.001x (mg/l) | 0.01x (mg/l) | 0.1x (mg/l) | 1x Environmental concentration (mg/l) | 10x (mg/l) | 100x (mg/l) | 1000x (mg/l) |
|------------------|---------------|--------------|-------------|---------------------------------------|------------|-------------|--------------|
| <b>Copper</b>    | 0.0005        | 0.005        | 0.05        | 0.5                                   | 5          | -           | -            |
| <b>Iron</b>      | 0.003         | 0.03         | 0.3         | 3                                     | 30         | -           | -            |
| <b>Manganese</b> | 0.002         | 0.02         | 0.2         | 2                                     | 20         | 200         | -            |
| <b>Zinc</b>      | -             | -            | 0.01        | 0.1                                   | 1          | 10          | 100          |
| <b>Nickel</b>    | 0.0005        | 0.005        | 0.05        | 0.5                                   | 5          | 50          | -            |



Table 3: Hydra morphology score table used to assess acute toxicity, based on the Wilby morphology score (Wilby, 1988).

|    |   |       |
|----|---|-------|
| 10 | Healthy, long tentacles and body reactive     |       |
| 9  | Partially contracted, slow reactions          |       |
| 8  | Clubbed tentacles, body slightly contracted   | Alive |
| 7  | Shortened tentacles, body slightly contracted |       |
| 6  | Tentacles and body shortened                  |       |
| 5  | Totally contracted, tentacles visible         |       |
| 4  | Totally contracted, no visible tentacles      |       |
| 3  | Expanded, tentacles visible                   | Dead  |
| 2  | Expanded, no visible tentacles                |       |
| 1  | Dead but intact                               |       |
| 0  | Disintegrated                                 |       |

Table 4: LC<sub>50</sub>, LOEC and NOEC values based on morphology for *Hydra attenuata* exposed to heavy metals individually and as a mixture. Toxicity Threshold (TT=(NOECxLOEC)/2). Actual concentrations measured in the environment are also presented.

| Metal     | Concentration in environment (mg/l) | LC <sub>50</sub> (mg/l) | LOEC (mg/l) | NOEC (mg/l) | TT (mg/l) |
|-----------|-------------------------------------|-------------------------|-------------|-------------|-----------|
| Mixture   | 1x                                  | 0.045x                  | 0.01x       | 0.001x      | 0.000005x |
| Copper    | 0.5                                 | 0.0225                  | 0.05        | 0.005       | 0.000125  |
| Iron      | 3                                   | 0.135                   | 0.03        | 0.003       | 0.000045  |
| Manganese | 2                                   | 20                      | 2           | 0.2         | 0.2       |
| Zinc      | 0.1                                 | 31.622                  | 10          | 1           | 5         |
| Nickel    | 0.5                                 | 2.25                    | 0.5         | 0.05        | 0.0125    |

Table 5: Pearson correlation coefficient of heavy metal pollutants and attachment, hydranth number and feeding behaviour endpoints.

|              | Mixture | Cu      | Fe      | Mn               | Ni      | Zn                |
|--------------|---------|---------|---------|------------------|---------|-------------------|
| Attachment   | -0.6742 | -0.5048 | -0.4882 | -0.6132          | -0.6666 | -0.6662           |
| Hydranth no. | -0.702  | -0.5032 | -0.5032 | <b>-0.9257**</b> | -0.717  | <b>-0.9996***</b> |
| Feeding      | -0.7013 | -0.5048 | -0.5048 | <b>-0.9308**</b> | -0.6848 | <b>-0.922**</b>   |

Significant results indicated by bold with significance set at \*p < 0.05, \*\*p < 0.005, \*\*\*p < 0.001

Table 6: A comparison of the heavy metal concentrations found from the Clyde, Thames and Ganga rivers with the maximum acceptable limits based on EU / UK legislative requirements.

| <b>Metal</b>     | <b>EU / UK<sup>a</sup><br/>(mg/l)</b> | <b>Average measured<br/>Concentration (mg/l)</b> | <b>River</b>                            |
|------------------|---------------------------------------|--|---|
| <b>Copper</b>    | 0.00376                               | 0.5  | Clyde, Glasgow, Scotland <sup>b</sup>   |
|                  |                                       | 0.001  | Thames, London, Canada <sup>c</sup>     |
|                  |                                       | 0.022  | Ganga, Kachlabridge, India <sup>d</sup> |
| <b>Iron</b>      | 1                                     | 3  | Clyde, Glasgow, Scotland <sup>b</sup>   |
|                  |                                       | 0.044  | Thames, London, Canada <sup>c</sup>     |
|                  |                                       | 0.0004   | Ganga, Kachlabridge, India <sup>d</sup> |
| <b>Manganese</b> | -                                     | 2  | Clyde, Glasgow, Scotland <sup>b</sup>   |
|                  |                                       | 0.011  | Thames, London, Canada <sup>c</sup>     |
|                  |                                       | -  | Ganga, Kachlabridge, India <sup>d</sup> |
| <b>Zinc</b>      | 0.0079                                | 0.1  | Clyde, Glasgow, Scotland <sup>b</sup>   |
|                  |                                       | 0.002  | Thames, London, Canada <sup>c</sup>     |
|                  |                                       | 0.00009  | Ganga, Kachlabridge, India <sup>d</sup> |
| <b>Nickel</b>    | 0.0086                                | 0.5  | Clyde, Glasgow, Scotland <sup>b</sup>   |
|                  |                                       | 0.004  | Thames, London, Canada <sup>c</sup>     |
|                  |                                       | 0.006  | Ganga, Kachlabridge, India <sup>d</sup> |

<sup>a</sup> SEPA (2018)

<sup>b</sup> Present study

<sup>c</sup> Environment and Engineering Services (2018)

<sup>d</sup> Central Water Commission (2018)

Table 7: Comparison of LC<sub>50</sub> for *H. attenuata* with those from other species for selected heavy metals.

| <b>Metal</b>     | <b>Organism</b>                 | <b>LC<sub>50</sub> (mg/l)</b> | <b>Source</b>                          |
|------------------|---------------------------------|-------------------------------|--|
| <b>Copper</b>    | <i>Hydra attenuata</i>          | 0.0225                        | Present study                          |
|                  | <i>Danio rerio</i>              | 0.01166                       | Alsop & Wood (2011)                    |
|                  | <i>Rasbora sumatrana</i>        | 0.0056                        | Shuhaimi-Othman <i>et al.</i> , (2010) |
|                  | <i>Capoeta fusca</i>            | 1.1                           | Ebrahimpour <i>et al.</i> , (2010)     |
|                  | <i>Poecilia reticulata</i>      | 0.0379                        | Shuhaimi-Othman <i>et al.</i> , (2010) |
| <b>Iron</b>      | <i>Hydra attenuata</i>          | 0.135                         | Present study                          |
|                  | <i>Daphnia magna</i>            | 0.23                          | García <i>et al.</i> , (2011)          |
|                  | <i>Salmo trutta</i>             | 0.05                          | Dalzell and MacFarlane (1999)          |
|                  | <i>Hyaella azteca</i>           | >1                            | Borgmann <i>et al.</i> , (2005)        |
| <b>Manganese</b> | <i>Hydra attenuata</i>          | 20                            | Present study                          |
|                  | <i>Rutilus rutilus caspicus</i> | 300                           | Hoseini <i>et al.</i> , (2014)         |
|                  | <i>Mogurnda mogurnda</i>        | 240                           | Harford <i>et al.</i> , (2015)         |
|                  | <i>Ceriodaphnia dubia</i>       | 6.2                           | Lasier <i>et al.</i> , (2000)          |
|                  | <i>Garra gotyla gotyla</i>      | 3.2                           | Sharma & Langer (2014)                 |
| <b>Zinc</b>      | <i>Hydra attenuata</i>          | 31.622                        | Present study                          |
|                  | <i>Danio rerio</i>              | 2.535                         | Alsop & Wood (2011)                    |
|                  | <i>Daphnia magna</i>            | 0.76                          | Lopes <i>et al.</i> , (2014)           |
|                  | <i>Capoeta fusca</i>            | 13.7                          | Ebrahimpour <i>et al.</i> , (2010)     |
|                  | <i>Lecane quadridentata</i>     | 0.12                          | Torres Guzman <i>et al.</i> , (2010)   |
| <b>Nickel</b>    | <i>Hydra attenuata</i>          | 2.25                          | Present study                          |
|                  | <i>Clarias gariepinus</i>       | 8.87                          | Ololade & Oginni (2010)                |
|                  | <i>Hyaella azteca</i>           | 2                             | Liber <i>et al.</i> , (2011)           |
|                  | <i>Danio rerio</i>              | 0.5898                        | Alsop & Wood (2011)                    |
|                  | <i>Chironomus dilutus</i>       | 119.5                         | Liber <i>et al.</i> , (2011)           |